

# Why can't my Beef Cows get Pregnant when Grazing Endophyte-infected Tall Fescue Pastures?

F. N. Schrick, G. M. Schuenemann, J. C. Waller, F. M. Hopkins and J. L. Edwards  
*Department of Animal Science, The University of Tennessee, Knoxville*

## **Introduction**

Tall fescue, a cool-season perennial grass, is one of the most commonly grown forages for over 8.5 million cattle in the United States [Hoveland, 1993]. Cattle suffering from fescue toxicosis experience decreased feed intake and performance, elevated respiration rate and body temperature, rough hair coats and necrosis of the extremities (tail, hooves, and ears) due to loss of circulation [Paterson et al., 1995]. Endocrine and reproductive effects of fescue toxicosis in cattle include decreased calving rate [Porter and Thompson, 1992] and pregnancy rates [Gay et al., 1988; Brown et al., 1992; Seals et al., 2005], reduced circulating concentrations of hormones such as cortisol, prolactin (PRL; [Porter, 1995]), progesterone (P<sub>4</sub>; [Mahmood et al., 1994; Jones et al., 2003]), and LH [Porter and Thompson, 1992; Mahmood et al., 1994]. This toxicosis results in estimated losses to the United States beef industry of \$609 million annually due to lowered conception rates and depressed body weight gains [Hoveland, 1993; Paterson et al., 1995].

We have made considerable progress in “narrowing the window” in determining the timing of reproductive loss associated with grazing infected tall fescue. We also know that addition of clover to our pastures will help reproductive performance in cattle, as does the addition of supplemental feeding of grain (etc.). This supplementation with clovers and grain is thought to have a “diluting” effect on the toxic component of tall fescue grass. In technical terms, the assumed “bad” component of tall fescue infected with the endophyte, *Neotyphodium coenophialum*, is an alkaloid known as “ergovaline” (produce by the endophyte) that negatively affects performance of the animal but plays a beneficial role on the hardiness of the grass [Hill et al., 1991; West et al., 1993; Ball et al., 1996; Thompson et al., 1999]. However, recent research by Hill et al. [2001] indicates that transport of the ergopeptine alkaloid “ergovaline” across ruminal gastric tissue is low as compared to the simple ergoline alkaloids lysergic acid and lysergol; thus, suggesting other alkaloids may play a larger role in tall fescue toxicosis.

So with all this said, the tall fescue research team at the University of Tennessee has focused their research attention on determining “how” and “when” the grazing of endophyte-infected tall fescue (E+) affects reproduction in cows and bulls. We performed these studies by either grazing tall fescue pastures (E+ or MaxQ, non-toxic endophyte, NTE) or by using a synthetic compound called ergotamine tartrate (referred to as ERGOT) to simulate the negative affects of ergovaline since ERGOT was commercially available, presented the same signs of tall fescue toxicosis, and we could control the nutritional status of the animal (thus removing nutrition from the equation as it relates to reproduction). Now to the reproduction part, we first wanted to know **when** consumption of endophyte-infected tall fescue had a negative effect on reproduction. There are several different periods of concern when looking at reproduction in a beef cattle setting, so we broke these time periods into different stages beginning with (1) the effect on the bull, (2) late pregnancy losses in the cow, (3) losses due to hormonal changes before estrus (heat), and (4) embryonic or uterine losses immediately following estrus.

### ***Effects on the Bull***

Few studies have focused on the beef bull, as related to fescue toxicosis, that had sufficient numbers or replicates to draw conclusions. Studies in mice [Zavos et al., 1988] and dairy bulls [Evans et al. 1988] had suggested a detrimental effect (mice) or no effect (Holstein bull calves) when consuming tall fescue seed or hay, respectively. Alamer and Erickson [1990] reported that yearling beef bulls grazing E+ tall fescue contained fewer Sertoli cells, suggesting impaired testicular function.

Recently, we completed analysis of data from a three-year project completed at Highland Rim Experiment Station with 96 yearlings beef (Angus and Gelbvieh) bulls. Year 1 [Schuenemann et al., 2005a] of the experiment involved yearling bulls receiving a control diet of corn silage supplemented with soybean meal (n=8) or a treated diet consisting of corn silage supplemented with soybean meal and “ergotamine tartrate (n=8) for a period of 224 days (November through June). This study was performed to control for the detrimental effect of tall fescue consumption on nutrition or weight loss. Again, feeding of ERGOT allows for us to control for the “nutrition factor” by regulating feed consumption and enables us to focus on the effects of the alkaloid on male fertility. Years 2 and 3 [Schuenemann et al., 2005b] utilized two sets of yearling beef bulls to evaluate the effect of actually grazing (experimental period of 224 days) tall fescue not infected (E-; Jesup MaxQ; n=10/year) or infected with the endophyte (E+) with (n=10/year) or without clover (n=20/year). Body weights, blood samples, forage samples and rectal temperatures were collected every 2 weeks. Every 60 days, scrotal circumference was recorded and semen collected for evaluation of motility and morphology. Testicular core temperatures were measured immediately before semen collection at the beginning of May and the end of June each year. Semen was extended immediately following collection and returned to the laboratory for evaluation through our in vitro fertilization program to determine fertilization potential and subsequent embryo development.

In brief, bulls consuming the diet supplemented with ergotamine tartrate had similar weight gains to control bulls as desired for the study. Scrotal circumference and semen motility and morphology were similar between treated and control bulls but fertilization potential (cleavage) was reduced (Table 1) in ERGOT bulls compared to controls.

Table 1. Ability of sperm collected from bulls fed a control or ergotamine tartrate-supplemented diet to fertilize bovine oocytes [Schuenemann et al., 2005a].

| <b>TRT</b>     | <b>REP<br/>(n)</b> | <b>COC<br/>(n)</b> | <b>PZ<br/>(n)</b> | <b>Cleav<br/>(%)</b>  | <b>Blast<br/>(%)</b> |
|----------------|--------------------|--------------------|-------------------|-----------------------|----------------------|
| <b>CON</b>     | 2                  | 200                | 169               | 69.2±3.3 <sup>a</sup> | 22.2±3.1             |
| <b>ERGOT</b>   | 2                  | 200                | 143               | 51.1±3.3 <sup>b</sup> | 22.0±3.1             |
| <b>P-value</b> |                    |                    |                   | 0.001                 | 0.96                 |
| <b>Lab Con</b> | 2                  | 100                | 86                | 74.4                  | 43.3                 |

<sup>a, b</sup> Least squares means differ within a column.

Reps: total number of replications per bull (Replicate 1, May 5<sup>th</sup>; Replicate 2, June 28<sup>th</sup>).

COC: cumulus oocyte complexes.

Cleav: number of putative zygotes cleaved.

Blast: blastocyst; percentage of cleaved embryos developing to blastocyst.

Subsequent development of embryos that cleaved was similar between treatments. However, testicular core temperatures were reduced in ERGOT bulls (Figure 1) compared to controls even though rectal temperatures were elevated suggesting a vasoconstrictive effect of consuming ergotamine tartrate on the testis.




Figure 1. Scrotal thermography measured at the time of semen collection on May 5<sup>th</sup> and June 28<sup>th</sup>. Scrotal temperatures recorded immediately before semen collections were lower in bulls administered ergotamine tartrate (ERGOT) compared to control (CON) animals (<sup>a, b</sup> Least squares means differ within treatments;  $P < 0.05$ ).

Schuenemann et al., 2005a

During Years 2 and 3, yearling beef bulls grazing E+ pastures without clover performed poorly (average daily gain) but no differences were noted in scrotal development or semen quality (as evaluated during a breeding soundness exam) in May and late June compared to bulls grazing E- (NTE; MaxQ) pastures. However, the fertilization ability or potential (ability to cleave) of the semen was reduced (Table 2) in bulls grazing E+ pastures without clover compared to E-. Again, rectal temperatures were elevated but with a reduction in testicular core temperatures (as measured by thermography) in bulls grazing E+ tall fescue pastures. Addition of clovers to the E+ pastures improved gain performance but fertilization potential of semen was not determined.

Table 2. Ability of sperm collected from bulls grazing tall fescue pastures to fertilize bovine oocytes [Schuenemann et al., 2005b].

| Variables  | NTE                   | E+                      | Lab Control |
|------------|-----------------------|-------------------------|-------------|
| Rep (n/yr) | 2                     | 2                       | 2           |
| COC (n)    | 850                   | 873                     | 278         |
| Cleav (%)  | 84 ± 2.4 <sup>a</sup> | 73.5 ± 3.1 <sup>b</sup> | 81 ± 3.8    |
| Blast (%)  | 30.1 ± 4.7            | 32.4 ± 5.5              | 30 ± 5.5    |
| Nuclei (#) | 76.5 ± 4.5            | 72.9 ± 4.8              | 73.6 ± 5.2  |

<sup>a, b</sup> Least squares means differ within a row,  $P < 0.05$ .

COC: cumulus oocyte complexes.

Cleav: number of putative zygotes cleaved.

Blast: blastocyst; percentage of cleaved embryos developing to blastocyst.

Nuclei #: total number of cells in blastocyst after fixation and Hoechst staining

Future studies needed on the male side include determining the effect of E+ tall fescue on reproduction parameters of MATURE bulls and sperm defects associated with fescue toxicosis and reduced fertilization ability.

**Effects on the Cow**

The next observation was that late pregnancy losses *were not occurring* at a higher rate in cows and heifers grazing E+ tall fescue as compared to animals grazing endophyte-free pastures. Heifers grazing infected tall fescue that were confirmed 30 days pregnant by ultrasonography stayed pregnant at an acceptable rate AND that the reproductive loss had already occurred by 30 days [Waller et al., 2001]. Thus, we deleted (2) *late pregnancy losses* as a concern in cows grazing endophyte-infected tall fescue. Similar results have been reported by several laboratories suggesting that reproductive losses occur early during pregnancy.

Next, we wanted to determine if changes in reproductive hormones or the follicle that produced the egg (oocyte) that would result in pregnancy differed in animals receiving ergotamine tartrate (ERGOT) to simulate fescue toxicosis [Seals et al., 2005]. In brief, pregnancy rates determined at 30 days postbreeding were reduced in cattle receiving ERGOT without any changes in reproductive hormones (Figure 2) or alterations in growth of the ovulatory follicle. Contrasting results have been reported concerning follicular development and alterations in reproductive hormones [Burke et al., 2001a; Burke et al 2001b; Jones et al., 2004] with others in agreement [Fanning et al., 1992]. Thus, in our laboratory with ergotamine tartrate, effects of fescue toxicosis were evident on pregnancy but not on any reproductive events that occurred immediately before breeding.




Figure 2. Area of the corpus luteum overlaid with progesterone concentrations during the ultrasonography period [Seals et al., 2005].

Next, we asked the question of whether the uterus was capable of maintaining a pregnancy when a good embryo was transferred into the reproductive tract on day 7 after estrus. In short, was fescue toxicosis causing a problem within the uterus? Rahe et al. [1991] suggested that losses were occurring after d 7 of pregnancy (following embryo transfer) but

left several unanswered questions. We observed that pregnancy rates were acceptable following transfer of frozen-thawed embryos (collected from cows not exposed to E+ tall fescue) in both groups of heifers that served as either controls or those receiving ERGOT [Schuenemann et al., 2005c]. Even though rectal temperatures were elevated, pregnancy rates were 50% in ERGOT heifers following embryo transfer. Thus, we deleted the *uterine environment after day 7* as a factor in reduced pregnancy rates during fescue toxicosis in heifers receiving ergotamine tartrate.

So when is fescue toxicosis affecting reproduction in the cow? The next experiment provided us a “smaller time frame”. Heifers were allotted into their respective treatment groups: receiving or not receiving ERGOT to simulate fescue toxicosis [Schuenemann et al., 2005c]. Estrus was synchronized and heifers (non-FSH stimulated) inseminated with semen from the same bull. Seven days later, single embryo recoveries were performed and embryos graded (evaluated) for quality and development. Embryos from heifers receiving ERGOT were reduced in quality and development (Figure 3). In other words, they were poor compared to our control embryos. Furthermore, recovery rate tended to be reduced in heifers supplemented with ergotamine tartrate.




Figure 3. Development (percentage of embryos that had attained the compacted morula or blastocyst stage) of embryos recovered on day 7 after estrus in heifers consuming ergotamine tartrate to simulate tall fescue toxicosis or serving as controls [Schuenemann et al., 2005c].

### Summary

So what does this all mean? Briefly, fescue toxicosis affects either the growing oocytes (egg) or the early embryo while still in the oviduct on the female side. Add in the effects of tall fescue on the sperm and we can understand why fertility is reduced. Throw in elevated temperatures during the summer months (June and July) with little (or no) clover and we could see a “reproductive wreck”. What we can suggest in terms of management is that we can remove cows from fescue for 30 days before and after breeding and see no effect on pregnancy rates (is this practical?). More practical would be to have your cows calve early (if spring calving) and get them exposed to the bull before the hot summer months occur (the same with your heifers). We intend to perform additional studies to determine if we can manage our females differently around the time of breeding (additional feeding, change

pasture locations and types, etc.) to improve pregnancy rates when grazing infected tall fescue.

Another factor to keep in mind would include the possibility that you have selected, over time, animals that will tolerate E+ tall fescue. You have culled those heifers that did not grow well. You have sold those cows that would not get pregnant. Has your herd adapted to grazing tall fescue? What about the bull? Will a bull that has not been exposed to tall fescue in his lifetime work on your farm? All the animals used in the above studies had been born and raised on the farm where the research was performed and consumed E+ tall fescue (with clover) their entire life. So have your animals been selected for adaptation? I'll leave that question for the genetic researchers!

### ***Acknowledgments***

Authors wish to thank the faculty, research associates, graduate and undergraduate students that have assisted over the years collecting data on these different studies. Special thanks to the University of Tennessee Experiment Stations for their assistance, especially Highland Rim, Plateau and East Tennessee (Knoxville) Research and Education Centers.

### ***References***

- Alamer, M. A., and B. H. Erickson. 1990. Effect of fungus-infected fescue on testicular development and hormone secretion in the beef bull. *J. Anim. Sci.* 68 (Suppl. 1): 402.
- Ball, D. M., C. S. Hoveland, and G. D. Lacefield. 1996. *Southern Forage* (2nd Ed.). Potash and Phosphate Institute and the Foundation for Agronomic Research. Norcross, GA.
- Brown, M. A., L. M. Tharel, H. A. Brown Jr., J. R. Miesner, and W. G. Jackson. 1992. Reproductive performance on Angus and Brahman cows grazing common bermudagrass or endophyte-infected tall fescue. *Prof. Anim. Sci.* 8:58-65.
- Burke, J. M., R. W. Rorie, E. L. Piper, and W. G. Jackson. 2001a. Reproductive responses to grazing endophyte-infected tall fescue by postpartum beef cows. *Theriogenology* 56:357-369.
- Burke, J. M., D. E. Spiers, F. N. Kojima, G. A. Perry, B. E. Salfen, S. L. Wood, D. J. Patterson, M. F. Smith, M. C. Lucy, W. G. Jackson, and E. L. Piper. 2001b. Interaction of endophyte-infected fescue and heat stress on ovarian function in the beef heifer. *Biol. Reprod.* 65:260-268.
- Evans, K. L., P. M. Zavos, R. W. Hemken, and J. A. Jackson. 1988. Effects of feeding endophyte-infected (*Acremonium-Coenophialum*) Ky-31 fescue hay on the reproductive development of Holstein bulls. *Theriogenology* 30:169-179.
- Fanning, M. D., J. C. Spitzer, D. L. Cross, and F. N. Thompson. 1992. A preliminary study of growth, serum prolactin and reproductive performance of beef heifers grazing *Acremonium coenophialum* infected tall fescue. *Theriogenology* 38:275-384.
- Gay, N., J. A. Boling, R. Dew, and D. E. Miksch. 1988. Effects of endophyte-infected tall fescue on beef cow-calf performance. *Applied Agriculture Research* 3:182-186.
- Hill, N. S., D. P. Belesky, and W. C. Stringer. 1991. Competitiveness of tall fescue as influenced by *Acremonium coenophialum*. *Crop Sci* 31:185-190.
- Hill, N. S., F. N. Thompson, J. A. Stuedemann, G. W. Rottinghaus, H. J. Ju, D. L. Dawe, and E. E. Hiatt III. 2001. Ergot alkaloid transport across ruminant gastric tissues. *J. Anim. Sci.* 79:542-549.
- Hoveland, C. S. 1993. Economic importance of *Acremonium* endophytes. *Agriculture, Ecosystem, and Environment* 44:3-12.
- Jones, K. L., S. S. King, K. E. Griswold, D. Cazac, and D. L. Cross. 2003. Domperidone can ameliorate deleterious reproductive effects and reduced weight gain associated with fescue toxicosis in heifers. *J. Anim. Sci.* 81:2568-2574.

- Jones, K. L., S. S. King, and M. J. Iqbal. 2004. Endophyte-infected tall fescue diets alters gene expressions in heifers luteal tissue as revealed by interspecies microarray analysis. *Mol. Reprod. Dev.* 67:154-161.
- Mahmood, T., R. S. Ott, G. L. Foley, G. M. Zinn, D. J. Schaeffer, and D. J. Kesler. 1994. Growth and ovarian function of weanling and yearling heifers grazing endophyte-infected tall fescue pastures. *Theriogenology* 42:1149-1158.
- Paterson, J., C. Forcherio, B. Larson, M. Samford, and M. Kerley. 1995. The effects of fescue toxicosis on beef-cattle productivity. *J. Anim. Sci.* 73:889-898.
- Porter, J. K. 1995. Analysis of endophyte toxins: fescue and other grasses toxic to livestock. *J. Anim. Sci.* 73:871-880.
- Porter, J. K., and F. N. Thompson. 1992. Effects of fescue toxicosis on reproduction in livestock. *J. Anim. Sci.* 70:1594-1603.
- Rahe, C. H., S. P. Schmidt, J. L. Griffin, C. A. Maness, D. I. Bransby, D. A. Coleman, R. L. Carson, and D. A. Stringfellow. 1991. Embryo survival is reduced in heifers grazing Kentucky-31 tall fescue infected with *Acremonium coenophialum*. *J. Anim. Sci.* (Supple. 1) 69:407.
- Schuenemann, G. M., J. L. Edwards, M. D. Davis, H. E. Blackmon, F. N. Scenna, N. R. Rohrbach, A. M. Saxton, H. S. Adair, F. M. Hopkins, J. C. Waller, and F. N. Schrick. 2005a. Effects of administration of ergotamine tartrate on fertility of yearling beef bulls. *Theriogenology* 63:1407-1418.
- Schuenemann, G. M., J. L. Edwards, F. M. Hopkins, F. N. Scenna, J. C. Waller, J. W. Oliver, A. M. Saxton, and F. N. Schrick. 2005b. Fertility aspects in yearling beef bulls grazing endophyte-infected tall fescue pastures. *Reprod. Fert. Dev.* 17:479-486.
- Schuenemann, G. M., M. E. Hockett, J. L. Edwards, N. R. Rohrbach, K. F. Breuel, and F. N. Schrick. 2005c. Embryo development and survival in beef cattle administered ergotamine tartrate to simulate fescue toxicosis. *Reprod. Biol.* 5:137-150.
- Seals, R. C., G. M. Schuenemann, J. W. Lemaster, A. M. Saxton, J. C. Waller, and F. N. Schrick. 2005. Follicular dynamics in beef heifers consuming ergotamine tartrate as a model of endophyte-infected tall fescue consumption. *JAVA* 4:97-102.
- Thompson, F. N., J. A. Studemann, and H. S. Hill. 1999. Antiquality factors associated with alkaloids in eastern temperate pasture grasses. Joint mtg. of American Forage and Grassland Council and the Society for Range Management. I. A.-Q. F. i. R. a. P. Forages. Omaha, NE.
- Waller, J. C., F. N. Schrick, M. C. Dixon, A. E. Fisher, A. M. Saxton and H. A. Fribourg. 2001. Tall fescue based forage systems for developing beef replacement heifers. *J. Anim. Sci.* 79:458.
- West, C. P., E. Izekor, K. E. Turner, and A. A. Elmi. 1993. Endophyte effects on growth and persistence of tall fescue along a water-supply gradient. *Agron J* 85:264-270.
- Zavos, P. M., D. R. Varney, J. A. Jackson, R. W. Hemken, M. R. Seigel, and L. P. Bush. 1988. Lactation in mice fed endophyte-infected tall fescue seed. *Theriogenology* 30: 865-875.